



# RJ VISION Pvt. Ltd.

Pioneer in Coaching For AIPMT(NEET) | JEE | PFC | NTSE | KVPY | OLYMPIAD

Code

**A**

# JEE – 2



(JEE – 2 FULL COURSE SOLUTION)

TEST ID : 302

TIME : 3 HR MM : 360

This Booklet contains 2 pages

**Important Instructions :**

1. The Answer Sheet is inside this Test Booklet. When you are directed to open the Test Booklet, take out the Answer Sheet and Fill in the particulars on **side-1** and **side-2** carefully with **blue/black** ball point pen only.
2. The test is of **3 hours** duration and Test Booklet contains **90** questions. Each question carries **4** marks. For each correct response, the candidate will get **4** marks. For each incorrect response, **one mark** will be deducted from the total scores. The maximum marks are **360**. [As per New Pattern]
3. Use **Blue/Black Ball point pen only** for writing particulars on this page/markings responses.
4. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
5. **On completion of the test, the candidate must handover the Answer Sheet to the invigilator in the Room/Hall. The candidates are allowed to take away this Test Booklet with them.**
6. The CODE for this Booklet is **A**. Make sure that the CODE printed on **Side-2** of the Answer Sheet is the same as that on this Booklet. In case of discrepancy, the candidate should immediately report the matter to the Invigilator for replacement of both the Test Booklets and the Answer Sheets
7. The candidates should ensure that the Answer Sheet is not folded. Do not make any stray marks on the Answer Sheet. Do not write your roll no. anywhere else except in the specified space in the Test Booklet/Answer Sheet.
8. Use of white fluid for correction is **NOT** permissible on the Answer Sheet.
9. Each candidate, must show on demand his/her Admission Card to the Invigilator.
10. No candidate, without special permission of the Superintendent or Invigilator, would leave his/her seat.
11. The candidates should not leave the Examination Hall without handing over their Answer Sheet to the Invigilator on duty and sign the Attendance Sheet twice. Cases where a candidate has not signed the Attendance Sheet the second time will be deemed not to have handed over Answer Sheet and dealt with as an unfair means case.
12. Use of Electronic/Manual Calculator is prohibited.
13. The candidates are governed by all Rules and Regulations of the Board with regard to their conduct in the Examination Hall. All cases of unfair means will be dealt with as per Regulations of the Board.
14. No part of the Booklet and Answer Sheet shall be detached under any circumstances.
15. The candidates will write the Correct Test Booklet Code as given in the Test Booklet/Answer Sheet in the Attendance Sheet.

Do not open this Test Booklet until you are asked to do so.

Name of the Candidate (in Capitals) : \_\_\_\_\_

Roll Number : \_\_\_\_\_

School : \_\_\_\_\_

Centre of Examination (in Capitals) (Vasna / Karelibaug/ / Others) \_\_\_\_\_

Candidate's Signature : \_\_\_\_\_ Invigilator Signature : \_\_\_\_\_

**ANSWER KEY**

Q.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
A.	3	3	2	4	2	2	1	2	1	2	1	3	1	2	3	4	4	1	4	4
Q.	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
A.	3	4	4	3	4	3	4	2	4	2	3	4	2	4	1	4	4	1	4	4
Q.	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
A.	3	1	3	4	3	4	4	3	1	3	2	4	3	3	1	3	2	1	3	4
Q.	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
A.	2	3	1	2	3	1	1	1	4	2	4	3	1	3	3	3	3	2	2	1
Q.	81	82	83	84	85	86	87	88	89	90										
A.	1	3	3	2	3	4	2	2	4	3										

**HINT - SHEET**

1.  $\frac{eV}{T} = \frac{\text{energy}}{\text{temperature}} = K$  Boltzman constant

2.  $a = \frac{v^2 - u^2}{2s} = \frac{100 - 400}{2 \times 0.5} = -300$

3.  $y = x \tan \theta = \frac{1}{2} g \frac{x^2}{u^2 \cos^2 \theta}$

$$y = x \cdot 1 - \frac{1}{2} \cdot \frac{x^2}{1}$$

$\therefore \tan \theta = 1 \quad \theta = 45^\circ$

$$\frac{u^2 \cos^2 \theta}{g} = 1$$

$$u = \frac{\sqrt{g}}{\cos \theta} = \sqrt{2g}$$

$$T = \frac{2u \sin \theta}{g} = \frac{2\sqrt{2g} \cdot \frac{1}{\sqrt{2}}}{g} = \frac{2}{\sqrt{g}}$$

10.  $PV = \mu RT$


$$\frac{V}{T} = \frac{\mu R}{P}$$

$$\frac{35}{28} \left( \frac{R}{6} \right) = \left( \frac{35}{28} + \frac{M}{32} \right) \frac{R}{(6+3)}$$

$M = 20 \text{ kg Ans.}$

20. As flux  $\phi = BA$

$$d\phi = \frac{\mu_0 i}{2\pi x} (adx)$$



$$\begin{aligned} \text{Total flux } \phi &= \int_b^{b+a} \frac{\mu_0 i a}{2\pi x} dx = \frac{\mu_0 i a}{2\pi} [\log_e x]_b^{b+a} \\ &= \frac{\mu_0 i a}{2\pi} \log_e \left( \frac{a+b}{b} + 1 \right) \\ &= \frac{\mu_0 a i_0 \sin \cot}{2\pi} [\log_e (a/b + 1)] \end{aligned}$$

$$\begin{aligned} \text{induced emf } e &= \frac{d\phi}{dt} \\ &= \frac{\mu_0 a i_0}{2\pi} \log \left( \frac{a}{b} + 1 \right) \frac{d}{dt} (\sin \cot) \\ &= \frac{\mu_0 a i_0 \omega \cos \cot}{2\pi} \log \left( \frac{a}{b} + 1 \right) \end{aligned}$$

21. In choice (3)

$$\text{RHS} = A \cdot B + A \cdot C = A \cdot (B + C) \neq A + B \cdot C$$

Remaining all are correct

22. For lyman series,  $\frac{1}{\lambda} = R \left( \frac{1}{1^2} - \frac{1}{\infty} \right) = R \Rightarrow \lambda = \frac{1}{R}$   
for paschen series,

$$\begin{aligned} \frac{1}{\lambda'} &= R \left( \frac{1}{3^2} - \frac{1}{\infty} \right) = \frac{R}{9} \\ \Rightarrow \lambda' &= \frac{9}{R} = 9\lambda \end{aligned}$$

SPACE FOR ROUGH WORK

24. Slope of the line  $\tan \theta = \frac{T_v}{T_c} > 1 \Rightarrow \theta > 45^\circ$

25.  $\frac{d^2x}{dt^2} = -ax - \omega^2x$

$\omega = \sqrt{a} \Rightarrow \frac{2\pi}{T} = \sqrt{a} \Rightarrow T = \frac{2\pi}{\sqrt{a}}$

29. Impedance  $Z = \sqrt{R^2 + X^2} = \sqrt{(30)^2 + (L\omega)^2}$   
 $= \sqrt{(30)^2 + \left(\frac{0.4}{\pi} \times 2\pi \times 50\right)^2}$   
 $= \sqrt{(30)^2 + (40)^2} = 50 \Omega$

Current (अवधि)  $i = \frac{V}{Z} = \frac{200}{50} = 4A$

30. at  $t = 4s$ ,  $R = \frac{R_0}{2^{t/T_0}} = 1600 \dots (1)$


at  $t = 8s$ ,  $R' = \frac{R_0}{2^{8/T_0}} = 100 \dots (2)$

$\therefore$  eq (1)/(2)  $\frac{R}{R'} = \frac{R_0}{2^{t/T_0}} \times \frac{2^{8/T_0}}{R_0} = \frac{1600}{100}$   
 $\Rightarrow 2^{(8-t)/T_0} = 16 \Rightarrow 2^{4/T_0} = 2^4 \Rightarrow 4/T_0 = 4$   
 $\Rightarrow T_0 = 1s$

from eq. (1)  $1600 = \frac{R_0}{2^4} \Rightarrow R_0 = 25600$


$\therefore$  at  $t = 6$

$R = \frac{R_0}{2^{6/T_0}} = \frac{25600}{2^6} = \frac{25600}{64} = 400$



$K_p = \frac{[NH_3]^2}{[N_2][H_2]^3} = \frac{(1/4)^2}{(0.5/4)(0.5/4)^3} = 256$

Equilibrium constant for the reaction



$K_c' = \frac{1}{\sqrt{K_c}} = \frac{1}{\sqrt{256}} = \frac{1}{16}$



$S = \sqrt{K_{sp}} = \sqrt{6.4 \times 10^{-5}} = 8 \times 10^{-3} \text{ mol/litre}$

$\therefore$  1 litre =  $10^{-3} \text{ m}^3$

$S = 8 \times 10^{-3} \text{ mol} / 10^{-3} \text{ m}^3 = 8 \text{ mol/m}^3$

61.  $R = \{(1, 3)(1, 5)(2, 3)(2, 5)(3, 5)(4, 5)\}$   
 $R^{-1} = \{(3, 1)(5, 1)(3, 2)(5, 2)(5, 3)(5, 4)\}$   
 $R \circ R^{-1} = \{(3, 3)(5, 5)(3, 5)(5, 3)\}$

62.  $\log_{(x)}\left(\frac{[x]}{x}\right)$  is defined ; if  $\frac{[x]}{x} > 0$ ,  $[x] > 0$  &

$[x] \neq 1$   
 $\Rightarrow x > 0$ ,  $x \in [1, \infty)$  and  $x \notin [1, 2) \Rightarrow x \in [2, \infty)$   
 For  $x \in [2, \infty)$ , we find that

$\log_{(x)}\left(\frac{[x]}{x}\right) = \log_{(x)}1 = 0$

$\therefore f(x) = \cos^{-1}(0) = \frac{\pi}{2}$  for all  $x \in [2, \infty)$ .

Hence, domain (f) =  $[2, \infty)$ , and range (f) =  $\left\{\frac{\pi}{2}\right\}$

74. Applying  $C_1 \rightarrow C_1 - (C_2 + C_3)$

$\begin{vmatrix} \sin x & \sin 2x & \sin 3x \\ 0 & 3 & 4 \sin x \\ 0 & \sin x & 1 \end{vmatrix}$   
 $= \sin x(3 - 4 \sin^2 x) - 3 \sin x - 4 \sin^3 x = \sin 3x$

$\therefore \int_0^{\pi/2} \sin 3x \, dx = \frac{1}{3}$

75.  $\int_{-\pi/2}^{\pi/2} \sin(x) \, dx$

$= 3 \int_0^1 \sin x \, dx$

$= 3[-\cos x]_0^1$   
 $= 3(1 - \cos 1)$


76. Eq<sup>n</sup> of tangent

$y - x(b+2) + (b+1) = 0$

Area of  $\Delta$  is

$= \frac{1}{2} \left[ \frac{b+1}{b+2} \times (b+1) \right] = 2$

$\frac{(b+1)^2}{b+2} = 4 \Rightarrow b = -3$



85. Obviously  $f(x) = \begin{cases} x^2, & x \leq 1 \\ 1, & x > 1 \end{cases}$

$\therefore f(x) = 1 = f(1)$ , so  $f(x)$  is continuous at  $x = 1$  and as such  $f(x)$  is continuous  $\forall x \in \mathbb{R}$ .

Further we note that  $f'(1-0) = 3$

and  $f'(1+0) = 0$

$\Rightarrow f(x)$  is not differentiable at  $x = 1$ .

Also  $f(x)$  exists  $\forall x \in \mathbb{R}$ ,  $x \neq 1$ .

Hence (3) are correct

SPACE FOR ROUGH WORK