

Translation :->

- * Process of polymerization of amino acids to form a polypeptide refers to translation.
- * The order and sequence of amino acids are defined by the sequence of bases in the m-RNA.
- * The amino acids are joined by a bond which is called peptide bond. formation of peptide bond requires energy **
Therefore,
In the first phase itself amino acids are activated in the presence of ATP and linked to their cognate t-RNA - a process commonly called as charging of t-RNA or aminoacylation of t-RNA to be more specific. If two such charged t-RNAs are brought close enough, the formation of peptide bond between them would be favoured energetically. The presence of a catalyst would enhance the rate of peptide bond formation.

- * The cellular factory responsible for synthesising proteins is the ribosome. The ribosome consists of structural RNAs and about 80 different proteins.
- * In its inactive state, it exists as two subunits; a larger and a smaller subunit.
- * When the small sub-unit encounters an m-RNA, the process of translation of the m-RNA to protein begins.
- * There are two sites in the larger subunit, for subsequent amino acids to bind to and thus, be close enough to each other for the formation of a peptide bond.
- * The ribosome also act as a catalyst (23S rRNA in bacteria is the enzyme-ribosome) for the formation of peptide bond.

- * A translational unit in m-RNA is the sequence of RNA that is flanked by the start codon (AUG) and stop codon and codes for polypeptide.
- * An m-RNA also has some additional sequences that are not translated and referred as untranslated regions (UTR). The UTRs are not at both 5'-end (before start codon) and at 3'-end (after stop codon). They are required for efficient translation process.

- * For initiation, the ribosomes binds to mRNA at the start codon (AUG) that is recognised by the initiator t-RNA.

- * The ribosomes proceed to the elongation phase of protein synthesis, during this stage, complexes composed of amino acid linked to t-RNA, sequentially bind to the appropriate codon in mRNA by forming complementary base pairs with the t-RNA anticodon.
- * The ribosome moves from codon to codon along the m-RNA.


- * Amino acids are added one by one, translated into polypeptide sequences dictated by DNA and represented by mRNA.

- * At the end, a release factor binds to the stop codon, terminating translation and releasing the complete polypeptide from the ribosome.

Translation

"m-RNA to polypeptide"


1. Activation of amino acids:-



* This process is called charging of t-RNA or aminoacylation of t-RNA

(2) Initiation of translation :-> It requires factors called initiation factors (IF1, IF2, IF3)


Eukaryotes have 9 initiation factors.



- * The initiating methionine accepting t-RNA is charged with non-formylated methionine ($t-RNA^{Met}$) in the cytoplasm of eukaryotes & formylated methionine ($t-RNA^{fMet}$) in prokaryotes, plastids and mitochondria.

- * t-RNA engaged in transferring formylated methionine is different than the one that transfers non-formylated methionine

(3) Elongation :- (Polypeptide chain formation)



3 step process

- (1) Codon recognition $\rightarrow GTP \rightarrow GDP + p ip$
- (2) Peptide bond formation \rightarrow (23S-rRNA = ribozyme)
- (3) Translocation $\rightarrow GTP \rightarrow GDP + p ip$ (ribosome moves over m-RNA)

elongation cycle takes less than a tenth of a second in bacteria and is repeated as each amino acid is added until the polypeptide is completed.

(4) Termination of Translation :-> This is final stage of translation process

